

Fertilgold® GREENS

INSTRUCTIONS

- FERTILGOLD® GREENS should be applied when tissue and soil analysis identifies the need and proper rate of application for this product.
- Best results will be obtained when application is concentrated in the active root zone.
- Can be applied in combination with compatible plant growth regulators, pesticides, or other granular fertilizers except urea.
- Apply FERTILGOLD® GREENS by broadcast, using airplane or ground machines.
- Application timing, intervals, and rates may vary according to individual crop requirement, stage of development, available nutrient levels in the soil, and overall nutritional status of the crop.
- Soil moisture is required for maximum bioactivity. If soil is dry, moisture should be provided by irrigation.
- Consult your local FERTILGOLD® Representative or other agricultural specialist for crop-specific recommendations.

This product is intended as a supplement to a complete fertilizer program and will not, by itself, provide the nutrients normally required by agricultural crops.

STORAGE AND DISPOSAL

Keep product in original bag. Do not transfer into food or drink containers. Always dispose of bag in accordance with local, state, and/or federal regulations.

SAFETY

KEEP OUT OF REACH OF CHILDREN

Dark reddish brown to black particles and dust having no odor. Ingestion of this product may cause gastrointestinal irritation or pain.

CONDITIONS OF SALE

The information contained on this label is believed to be accurate and reliable. Buyer and user acknowledge and assume all liability resulting from the use of this material. Follow directions carefully. Timing, method of application, weather, crop conditions, and other factors are beyond the control of the seller.

0-0-0

GUARANTEED ANALYSIS

ALSO CONTAINS NON-PLANT FOOD INGREDIENTS:

Total Humic Acid (HA) and

Fulvic Acid (FA).....60.0%-70.0%*

*Determined using V&B (Verploegh and Brandvoid) Method

Derived from: Oxidized Leonardite

Designed to be applied to the soil.

Avoid spreading during high humidity, or mixing with high moisture fertilizers such as urea.

Applications can be made as often as every 30 days as needed.

APPLICATION

METHOD	SUGGESTED RATE PER ACRE Field Crops / Tree or Vine Crops
Soil banded or side dressed	Up to 75 lbs / up to 150 lbs
Soil broadcast incorporated	Up to 150 lbs / up to 300 lbs

Net Wt. 2,000 lb (1 Ton)
Net Wt. 907 kg
Lot No. 12345678

© 2017–2018 *Registered Trademark of and
Produced by: Bio Huma Netics, Inc.
1331 W. Houston Ave., Gilbert, AZ 85233
480.961.1220
FGO-640-08212018

Fertilgold.com

