

# Fertilgold<sup>®</sup> Mg

## INSTRUCTIONS

- Designed for both soil and foliar application.
- FERTILGOLD<sup>®</sup> Mg should be applied when tissue and soil analysis identifies the need and proper rate of application for this product.
- Best results will be obtained when application is concentrated in the active root zone or on the leaf surface.
- Can be applied in combination with compatible plant growth regulators, pesticides, or other liquid fertilizers. If compatibility is in question, jar test a small quantity.
- Apply foliar sprays with sufficient water to ensure uniform coverage without running off leaf surfaces. For maximum results, apply in the early morning or in the evening when temperatures are optimum for foliar application.
- Do not apply FERTILGOLD<sup>®</sup> nutrients during the hottest time of the day or to plants under drought or flood stress.
- Application timing, intervals, and rates may vary according to individual crop requirement, stage of development, available nutrient levels in the soil, and overall nutritional status of the crop.
- Consult your local FERTILGOLD<sup>®</sup> Representative or other agricultural specialist for crop-specific recommendations.

This product is intended as a supplement to a complete fertilizer program and will not, by itself, provide all of the nutrients normally required by agricultural crops.

## STORAGE AND DISPOSAL

Keep product in original container. Do not transfer into food or drink containers. Triple rinse when empty for recycling. Always dispose of container in accordance with local, state, and/or federal regulations. Do not store this product below 50°F (10°C) or above 90°F (30°C).

© 2017–2018 \*Registered Trademark of and Produced by: Bio Huma Netics, Inc.  
1331 W. Houston Ave., Gilbert, AZ 85233  
480.961.1220  
FGO-880-08212018

0-0-0

## GUARANTEED ANALYSIS

Sulfur (S) ..... 6.0%  
Magnesium (Mg) ..... 5.0%

Derived from:  
Magnesium Sulfate

ALSO CONTAINS NON-PLANT FOOD INGREDIENT:  
3.0% Organic Matter (derived from leonardite)

SHAKE WELL BEFORE USE.

Do not apply this product foliarly in concentrations greater than 5%.

Applications can be made as often as every 7 to 10 days as needed.

## APPLICATION

METHOD	SUGGESTED RATE PER ACRE Field Crops / Tree or Vine Crops
Foliar band application at 50% coverage	Up to 2 quarts / –
Foliar broadcast or sprinklers: solid, set, linear, pivot (100% speed)	Up to 1 gallon / up to 2 gallons
Soil banded or injected, through drip tape or microsprinklers	Up to 2 gallons / up to 4 gallons
Soil broadcast spray incorporated, flood, or furrow irrigated	Up to 4 gallons / up to 8 gallons

pH: 6.0 - 7.0

10.43 lb/gal @ 68°F  
1.25 kg/L @ 20°C

## SAFETY

KEEP OUT OF REACH OF CHILDREN

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.html>.

*A slightly hazy amber liquid having a characteristic odor. The liquid and mists can cause eye irritation and may cause slight to moderate skin irritation. Inhalation of aerosols or mists may cause irritation to the respiratory tract. Ingestion may cause irritation to the entire gastrointestinal tract with central nervous system depression, lack of reflexes, sweating, flushing, decreased blood pressure and hypocalcemia.*

CLASSIFICATION: HAZARD CATEGORY 3 - MILD SKIN IRRITATION

SIGNAL WORD: WARNING

HAZARD STATEMENT: H316 - WARNING – causes mild skin irritation

PRECAUTIONARY STATEMENT: P332+P313; If skin irritation occurs: Get medical attention/advice.

## CONDITIONS OF SALE

The information contained on this label is believed to be accurate and reliable. Buyer and user acknowledge and assume all liability resulting from the use of this material. Follow directions carefully. Timing, method of application, weather, crop conditions, and other factors are beyond the control of the seller.

275 gallons • Net Wt. 2,868 lb  
1,041.15 liters • Net Wt. 1,301 kg  
Lot No. 12345678

Fertilgold.com

